

CastAway DGTM

TEA SEED MEAL FERTILIZER

Fall Best Management Practice (BMP) Guidelines

Fall weather is just around the corner and turf managers need to plan ahead for the unique needs of the season. Now is the time to stock up on CastAway DG because the window of correct usage is time-sensitive once earthworm activity resumes as rains return and temperatures fall. With that in mind, here are some fall Best Management Practice (BMP) application guidelines for CastAway DG.

- ▶ Apply at first notice of earthworm casting occurrence
- ▶ First application should be at the labeled HIGH rate (6.0 lbs./M, covers 8,000 ft²).
- ▶ Sequential applications can be at the labeled LOW (3.0 lbs./M, covers 13,300 ft²) or HIGH rate .
- ▶ Reapplication interval can be highly variable, dependent on soil and weather conditions
- ▶ Reapply at first observed reoccurrence of casting mounds
- ▶ Multiple applications are likely necessary for complete fall control
- ▶ Water in all applications, beginning with 0.10" of water, followed by a second application if necessary to fully disperse product
- ▶ Earthworms may relocate; treat putting green edges if treating collars to prevent casting mounds on greens
- ▶ Uniform application is best achieved by using The Andersons Model 2000 or 2000 SR spreaders, cone setting 8



Natural fertilizer with high protein content



Nutrients derived from tea plant seed meal



Dispersing Granule (DG) Technology provides rapid incorporation



FOR MORE INFORMATION

Contact your local authorized distributor of The Andersons products, your Territory Manager from The Andersons, or call 800-253-5296.

AndersonsPlantNutrient.com

Twitter Facebook Instagram @TheAndersonsPro
YouTube AndersonsPro